
IMPACT OF YOGIC PRACTICES ON SELECTED BODY COMPOSITION MEASURES AND HIGH DENSITY LIPOPROTEINS AMONG OBESE WOMEN

S. Chidambara Raja

Associate Professor, Department of Physical Education and Sports Sciences, Annamalai University.

Abstract:-The purpose of the study was to find out the effect of yogic practices on selected body composition measures and high density lipoproteins among obese women. To achieve this purpose, 20 obese women, with BMI of 30-40 (W/H²), were randomly selected as subjects from various places around Chidambaram. The age of the subjects were ranged from 30 to 35 years. The subjects were further classified at random into two equal groups of 10 subjects each, in which, group - I underwent yogic practices for six days (Monday to Saturday) per week for sixteen weeks and group - II acted as control who were not allowed to attend any special training. The selected criterion variables such as percentage of body fat, body mass index and high density lipoproteins were measured before and after the yogic practice period. The selected criterion variables were assessed by using Deurenberg et al formula, Quetelet index and Boehringer Mannheim kit method. The collected data were statistically analysed by using Analysis of Covariance (ANCOVA). From the results of the study it was found that there was a significant reduction in percentage of body fat ($p > .05$) and body mass index ($p > .05$) and a significant increase in high density lipoprotein level ($p > .05$) after the yogic practice when compared with the control group. It was concluded from the result of the study, that yogic practice is a better tool to reduce the percentage of body fat and body mass index and increase the level of high density lipoprotein.

Keywords:Yogic Practices, Body Composition Measures, HDL.

INTRODUCTION

Obesity is a medical condition in which excess body fat has accumulated to the extent that it may have an adverse effect on health, leading to reduced life expectancy and/or increased health problems.[1] It is a metabolic disorder which is affecting the people throughout the world and commonly caused by a combination of excessive food energy intake, lack of physical activity, genetic susceptibility, and other psychological problems, although a few cases are caused primarily by genes, endocrine disorders, medications or psychiatric illness.[2] The negative health (obesity) consequences are less or more insulin resistance, chances of occurring type 2 diabetes, asthma, hypertension, increase in high total cholesterol, low density lipoproteins, triglycerides and lowering the high density lipoproteins in blood, become sleep apnea, attaining early puberty, etc.[3] Indexes associated with high risk in obese persons often return to normal with appropriate physical activities, dietary habits, and a small weight loss even when body weight and percentage body fat remain above recommended amounts.[4]

Women suffer a disproportionate burden of disease attribute to overweight and obesity when comparing with men and the women those who are measuring more than 30 percent of the ideal body weight are becoming obese.[5] Women with body weight measuring up to 30 percent more than that of the ideal are known as Obese. The body mass index (BMI), is the most popular and effective way of calculating Female Obesity.[5]

Please cite this Article as :S. Chidambara Raja, "IMPACT OF YOGIC PRACTICES ON SELECTED BODY COMPOSITION MEASURES AND HIGH DENSITY LIPOPROTEINS AMONG OBESE WOMEN" : Academic Sports Scholar (Nov ; 2014)

Those children who have BMI of above 95% percentiles are in obese.[6] More children aged 2 to 5 years are obese, as are 17 percent of children aged 6 to 19 according to the Centers of Disease Control and Prevention (CDC).[7] The primary problems for obese children are psychological or emotional.[8] It is also evident that increasing mortality rate during adolescent are due to childhood obesity.[9] A 2008 study has found that children who are obese have carotid arteries which have prematurely aged by as much as thirty years as well as abnormal levels of cholesterol.[10] The obese children were abused and teased by their same age group[11] and also by their family members quite often.[12][13]

Yoga is a spiritual science for the integrated and holistic development of physical, mental and spiritual aspects of our well being.[14] Yoga is originated in India many thousands of years ago and it is the oldest system of personal development in the world, encompassing body, mind and spirit.[15,16] Yogsana have a deeper significant value in the development of the physical, mental and spiritual personality, whereas pure exercises only have a physical effect on the muscles and bones.[17] Yoga poses are also designed to tone and exercise the muscles of the body to eliminate excess fat, and make it more flexible and stronger.[18] Yogic practice reduces the obesity and also reduces the risk factors associated with obesity.[19] A study shows there was a significant reduction in total cholesterol and increase in HDL after twelve weeks of yoga practices.[20] Various researches suggest that yoga exercise improves the BMI of sedentary human beings including women.[21,22,23,24]

The exact body fat percentage cannot be precisely determined, but multiple methods are used to estimate it.[25] There is no single ideal percentage of body fat for everyone. Levels of body fat are epidemiologically dependent on sex and age.[26] There are many methods examine the percentage of body fat, such like, underwater body weight, skinfold test, bioelectrical impedance analysis, etc. The percentage of body fat will also be estimated with person's body mass index (BMI) by applying Deurenberg et al formula.[27] Body mass index (BMI) has recently gained favor as a better measure of adiposity.[28,29]

METHODOLOGY

The purpose of this study was to find out the effect of yogic practices on percentage of body fat, body mass index and high density lipoproteins among obese women. To achieve the purpose of the present study, 20 obese women with the BMI of 95 percentile[30] or above and who were studying in various schools around Chidambaram, Tamilnadu were randomly selected as subjects. The age of the subjects were ranged from 14 to 16 years (mean age = 15.1 ± 0.3 years). All the subjects were residing at their home, so, the food habits were not same and could not be measured. The selected subjects were divided into two equal groups of ten subjects each. Group - I considered as experimental group who underwent yogic practices for sixteen weeks, six days (Monday to Friday) per week on selected yogic exercises (appendix – I) and the same were taught by yoga teachers from School of Yoga Studies, Annamalai University, Annamalainagar, Chidambaram. Group - II considered as control that did not undergo any training programme or physical activity (either strenuous or recreational) throughout the experimental period. The data were collected on selected criterion variables such as percentage of body fat was assessed by using Deurenberg et al[27,31] formula, body mass index was assessed by Quetelet index[32] and high density lipoproteins was assessed by phosphotungstate/Mg²⁺ method, using the reagent from Boehringer Mannheim Lab, Germany[33] after taking 5 ml of blood from each subject by venous puncture method by the lab technicians, under the supervision of a qualified doctor, before and after the sixteen weeks of yogic practices as pre and post test. The blood samples were collected only after the consulting and getting permission from their parents. Analysis of covariance (ANCOVA) was applied to find out the significant difference if any between the experimental and control groups.

Table – II
Analysis of Covariance on Percentage of Body Fat Body, Body Mass Index and High Density Lipoproteins of Yogic Practice Group and Control Group

Variable Name	Group Name	Yogic Practice Group	Control Group	'F' Ratio
Percentage of Body Fat (in Percentage)	Pre-test Mean ± S.D	28.2521 ± 1.8236	29.6861 ± 1.893	2.991
	Post-test Mean ± S.D.	25.8926 ± 1.5406	29.8121 ± 2.126	18.91*
	Adj. Post-test Mean	25.561	29.299	32.351*
Body Mass Index (kg/m ²)	Pre-test Mean ± S.D	31.8682 ± 1.3317	31.1816 ± 1.2628	0.1962
	Post-test Mean ± S.D.	29.3839 ± 0.8623	31.2125 ± 1.452	19.253*
	Adj. Post-test Mean	27.896	31.531	29.632*
High Density Lipoproteins (mg/dl)	Pre-test Mean ± S.D	40.22 ± 2.278	39.67 ± 1.976	0.989
	Post-test Mean ± S.D.	42.56 ± 1.884	38.89 ± 2.176	5.387*
	Adj. Post-test Mean	42.198	38.773	18.551*

*Significant at 0.05 level of confidence.(The table values required for significance at 0.05 level of confidence for 1 and 18 & 1 and 17 are 4.41 and 4.45 respectively).

RESULTS

The collected data prior to and after the yoga practice on percentage of body fat, body mass index and high density lipoprotein were analyzed by applying Analysis of Covariance (ANCOVA) are presented in table – II. In all the cases, .05 level of confidence was fixed to test the significance, which was considered as an appropriate.

After applying the Analysis of Covariance, the result of this study shows that there was a significant decrease in percentage of body fat (Exp. Gr. Pre-mean = 28.2521 ± 1.8236 Vs Post-mean = 25.8926 ± 1.5406 & Cont. Gr. Pre-mean = 29.6861 ± 1.893 Vs. Post-test mean = 29.8121 ± 2.126) and body mass index (Exp. Gr. Pre-mean = 31.8682 ± 1.3317 Vs. Post-mean 29.3839 ± 0.8623 & Cont. Gr. Pre-mean = 31.1816 ± 1.2628 Vs. Post-mean 31.2125 ± 1.452) and also there was a significant increase in high density lipoproteins (Exp. Gr. Pre-mean = 40.22 ± 2.278 Vs. Post-mean 42.56 ± 1.884 & Cont. Gr. Pre-mean = 39.67 ± 1.976 Vs. Post-mean 38.89 ± 2.176) only for yogic practice group.

Further, comparing the adjusted post-test means of the criterion variables (between yogic practice group and control group), such as percentage of body fat (Exp. Gr. = 25.561 Vs. Cont. Gr. = 29.299 & F = 32.351, p < 0.05), body mass index (Exp. Gr. = 27.896 Vs. Cont. Gr. = 31.531 & F = 29.632, p < 0.05) the yogic practice group was significantly decreased when compared with control group. High density lipoprotein was significantly increased for yogic practice group (Exp. Gr. = 42.198 Vs. Cont. Gr. = 38.773, & F = 18.551, p < 0.05) with df 1,17.




Figure- I: Bar diagram showing the mean values of % of body fat between yogic practice group and control group





Figure- II: Bar diagram showing the mean values of BMI between yogic practice group and control group

Figure- III: Bar diagram showing the mean values of HDL between yogic practice group and control group



DISCUSSION

In the present study the subjects were obese children and their BMI was at 95th percentile.

- 1.The reduction in percentage of body fat and body mass index was significant for yogic practice group when compared with the control group.
- 2.There was a significant increase in high density lipoprotein cholesterol for yogic practice group when compared with the control group.

CONCLUSIONS

- 1.The results of the study revealed that there was a significant reduction in percentage of body fat after the yogic practice period. This result is in line with that of the study earlier conducted by Pal et al[34] and Shenbagavalli and Divya[35] found that there was a significant reduction in percentage of body fat after the yogic practice. Ruhall, Bhandari and Chakravarti[36] also found that there was a significant reduction in percentage of body fat after the pranayama practice.
- 2.The result of the study also shown that there was a significant reduction in body mass index (BMI) after the yogic practice period, when compared with the control group. The findings of Kumari et al[37], Dhananjai et al[38] and Chen et al[39] also found that there was a significant decrease in body mass index after the yogic practice period. Ankad et al[40] also found that there was a significant decrease in body mass index after the pranayama practice.
- 3.The result of the study shown that there was a significant increase in high density lipoproteins after the yogic practice period, when compared with the control group. The findings of Telles et al[41]

found that there was a significant increase in high density lipoproteins after the yogic practice period. 4. The overall result of the study shown that there was a significant reduction in percentage of body fat, body mass index and high density lipoproteins after the experimental period. In this study, no attempt was taken to control the diet. But, in future, if the effort will be taken, the reduction in percentage of body fat, body mass index and high density lipoproteins will be higher.

REFERENCE:

- [1] Haslam DW and James WP (2005). "Obesity". *Lancet*, 366:9492, 1197–209.
- [2] Retrieved from <http://en.wikipedia.org/wiki/Obesity> on 12/07/2014.
- [3] Retrieved from <http://www.obesity.org/resources-for/childhood-overweight.htm> on 15-07-2014.
- [4] Aherlthy RP and Black DR (1996). "Healthy Body Weights: an Alternative Perspective". *American Journal of Clinical Nutrition*, 63, 448–451.
- [5] Retrieved from <http://www.patientsmedical.com/women/obesity/> on 24-07-2014.
- [6] Retrieved from http://kidshealth.org/kid/grow/body_stuff/bmi.html on 12-07-2014.
- [7] Kimberley Wonderly, "What is a Normal BMI for Women", Retrieved from <http://www.livestrong.com/article/88045-normal-bmi-women/> on 2-06-2014.
- [8] Great Britain Parliament House of Commons Health Committee (May 2004). *Obesity - Volume 1 - HCP 23-I, Third Report of session 2003-04. Report, together with formal minutes*. London, UK: TSO (The Stationery Office). ISBN 978-0-215-01737-6. Retrieved 2007-12-17.
- [9] Must A, Jacques PF, Dallal GE, Bajema CJ, Dietz WH (November 1992). "Long-term morbidity and mortality of overweight adolescents. A follow-up of the Harvard Growth Study of 1922 to 1935". *The New England Journal of Medicine*, 327:19, 1350–5.
- [10] Facts for Families, "Obesity in Women Teens", Retrieved from http://www.aacap.org/App_Themes/AACAP/docs/facts_for_families/79_obesity_in_women_and_teens.pdf on 15-07-2014.
- [11] Janssen I, Craig WM, Boyce WF, Pickett W (2004). "Associations between overweight and obesity with bullying behaviors in school-aged women". *Pediatrics*, 113:(5), 1187–94.
- [12] Retrieved from <http://web.archive.org/web/20020602165343/http://obesity.org/discrimination/educa.shtml> on 12-07-2014.
- [13] Gurvinder Kalra, Avinash De Sousa, Sushma Sonavane and Nilesh Shah (January – June 2012). "Psychological Issues in Paediatric Obesity". *Indian Psychiatry Journal*, 21:2, 11-17.
- [14] Bhavanani, A.B. (2011). "Application of Yoga Concept in the Health Improvement". In: P.Nikic, ed. *Proceedings "Yoga – the Light of Microuniverse" of the International Interdisciplinary Scientific Conference "Yoga in Science – Future and Perspectives"*, September 23-24, 2010, Belgrade, Serbia. Belgrade: Yoga Federation of Serbia, p. 189-193.
- [15] Swami Vishnu Devananda, *The Sivananda Companion to Yoga*, (New York: Fireside Book, Simon and Schuster, 2000), p. 10.
- [16] Retrieved from <http://www.minddisorders.com/Py-Z/Yoga.html> on 24-04-2012.
- [17] Chidambara Raja (November – December, 2012). "Effect of Yogic Practices on Flexibility Cholesterol and Blood Pressure". *Online International Interdisciplinary Research Journal*, 2:6, 221-225.
- [18] Retrieved from <http://www.abc-of-yoga.com/yoga-and-health/yoga-for-women.asp> on 06-05-2012.
- [19] Dhananjai S, Sadashiv, Kumar Rajjan M.P.S. Negi and Dr. Sunita Tiwari (July 2011). "Effect of yoga practice in Management of Risk Factors with Obesity: A Pilot study". *Indians Streams Research Journal*, 1:4, 1-7.
- [20] Mathew Ted Ballwin, "Asthangha Yoga for Women – A Viable Alternative for Weight Management and Psychological Well Being", retrieved from <http://www.healthandyoga.com/html/news/therapy/yogakids.aspx> on 12-07-2014.
- [21] T.L. Chen, H.C. Mao, C.H. Lai, C.Y. Li and C.H. Kuo, "The Effect of yoga Exercise Intervention on Health Related Physical Fitness in School – Age Asthmatic Women". *Hu Li Za Zhi*, 56:2, (April 2009), 42-52.

- [22] Suchetha Kumari N, Damodara Gowda KM, Sukesh N, Madhu LN and Kathyayini (September 2011). "Effect of Yoga Therapy on Body Mass Index and Oxidative Status". *Nitte University Journal of Health Science*, 1:1-3, 10-14
- [23] Balamurugan D, Ashok Kumar R and Dr. Karikalan I (January to June 2013). "Effect of Varied Yogic Practices on Body Mass Index Component of Obese Engineering College Men Students". *Asian Journal of Physical Education and Computer Science in Sports*, 8:1, 43-44.
- [24] Yokesh TP and Chandrasekaran K. (2011). "Effect of Yogic Practice on Selected Physical Fitness among Overweighted School Women". *Recent Research in Science and Technology*, 3:9, 43-45.
- [25] Retrieved from <http://weightloss.about.com/od/glossary/g/percentbodyfat.htm> on 6-8-2012.
- [26] Jackson, AS; Stanforth, PR; Gagnon, J; Rankinen, T; Leon, AS; Rao, DC; Skinner, JS; Bouchard, C; Wilmore, JH (2002). "The Effect of Sex, Age and Race on Estimating Percentage of Body Fat from Body Mass Index: The Heritage Family Study". *International Journal of Obesity and Related Metabolic Disorder*, 26:6, 789-96.
- [27] Deurenber P, Westrate JA and Seidell JC (March 1991). "Body Mass Index as a Measure of Body Fatness: Age- and Sex-Specific Prediction Formulas". *British Journal of Nutrition*, 65:2, 105-14.
- [28] Kraemer H, Berkowitz RI and Hammer LD (1990). "Methodological Difficulties in Studies of Obesity, I: Measurement Issues". *Ann Behav Med*. 12, 112-118.
- [29] Must A, Dallal GE and Dietz WH (1991). "Reference Data for Obesity: 85th and 95th Percentiles of Body Mass Index (wt/ht²) and Triceps Skinfold Thickness". *American Journal of Clinical Nutrition*, 53, 839 - 846.
- [30] Retrieved from <http://www.webmd.com/parenting/raising-fit-kids/weight/kids-bmi-for-parents> on 22-06-2014.
- [31] Retrieved from <http://www.iarc.fr/en/publications/pdfs-online/prev/handbook6/Handbook6-1.pdf> on 22-6-2014.
- [32] Retrieved from http://en.wikipedia.org/wiki/Body_mass_index on 21-06-2014.
- [33] Pierre N.M. Demacker, Marja Hessels, Helga Toenhake-Dijkstra and Henk Baadenhuijsen, (April 1997). "Precipitation Methods for High Density Lipoprotein Cholesterol Measurement Compared, and Final Evaluation Under Routine Operating Conditions of a Method with a Low Sample-to-Reagent Ratio". *Clinical Chemistry*, 43:4, 663-668.
- [34] Pal A, Srivastava N, Tiwari S, Verma NS, Narain VS, Agrawal GG, Natu SM and Kumar K (June 2011). "Effect of Yogic Practices on Lipid Profile and Body Fat Composition in Patients of Coronary Artery Disease". *Complementary Therapies in Medicine*, 19:3, 122-7.
- [35] Shenbagavalli A and Divya K (2010). "The Effect of Specific Yogic Exercises and Combination of Specific Yogic Exercises with Autogenic Training on Selected Physiological Psychological and Biochemical Variables of College Men Students". *Journal of Exercise Science and Physiotherapy*, 6:2, (2010), 94-101.
- [36] Ajay Singh Ruhall, Rakesh Bhandari and Ranjan Chakravarti (2010). "Effect of Kapalabhati on Selected Body Composition Variables". *British Journal of Sports Medicine*, 44, i 70.
- [37] N. Suchetha Kumari, K.M. Damodara Gowda, N. Sukesh, L.N. Madhu and Kathyayani, "Effect of Yoga Therapy on Body Mass Index and Oxidative Status", *Nitte University Journal of Health Science*, 1.:1-3, (September 2011). 10-14.
- [38] S. Dhananjai, Sadashiv, Kumar Rajjan, M.P.S. Negi and Sunita Diwari, "Effect of Yoga Practice in the Management of Risk Factors Associated with Obesity: A Pilot Study", *Indian Streams Research Journal*, 1:6, (July 2011).
- [39] T.L. Chen, H.C. Mao, C.H. Lai, C.Y. Li and C.H. Kuo, "The Effect of Yoga Exercise Intervention on Health Related Physical Fitness in School-age Asthmatic Women", *Hu Li Za Zhi*, 56:2, (April 2009), 42 - 52.
- [40] Roopa B. Ankad, Anita Herur, Shailaja Patil, G.V. Shashikala and Surekharani Chinnagudi, "Effect of Short-term Pranayama and Meditation on Cardiovascular Functions in Healthy Individuals", *Heart Views*, 12:2, (April - June 2011), 58 - 62.
- [41] S. Telles, V.K. Naveen, A. Balakrishna and S. Kumar, "Short Term Health Impact of a Yoga and

Diet Change Program on Obesity”, Med. Sci. Monit., 16:1, (January 2010), 35-40.

FORMULAS:

Calculating the Percentage of Body Fat:

$$\text{Child Body Fat \%} = (1.51 \times \text{BMI}) - (0.70 \times \text{Age}) - (3.6 \times \text{gender}) + 1.4$$

Where as the gender: Male = 1, Female = 0. (Retrieved from <http://www.halls.md/bmi/fat.htm> on 15-08-2012).

Calculating Body Mass Index (BMI):

The weight in kilograms divided by the square of the height in meters, used in the assessment of underweight and obesity. (Retrieved from

<http://medical-dictionary.thefreedictionary.com/Quetelet+Index> on 19-05-2012).

Table - I
TRAINING SCHEDULE FOR YOGIC PRATICE GROUP

List of Yogasana & Pranayama	Weeks	Duration	Maintaining Duration	Recovery in between Yogasanas	Repetitions	Frequency	Warming up and cooling down
Padmasana	1-3 Weeks	20 min.	30 seconds	30 seconds	2	Monday Tuesday Wednesday Thursday Friday & Saturday	5 – 10 Minutes
Trikonasana			30 seconds	30 seconds			
Dhanurasana			30 seconds	30 seconds			
Shashangasana			1 minute	30 seconds			
Patchimosthasan			1 minute	30 seconds			
Meditation–Omkar.			1 minutes	30 seconds			
Pranayama – Nadisuthi			1 minutes	30 seconds			
Shavasana			2 minutes		1		
As in previous week	4 – 6 Weeks	40 min	20 minutes		2	Monday Tuesday Wednesday Thursday Friday & Saturday	5 – 10 Minutes
Bhujangasana			1 minute	30 seconds			
Shalabasana			1 minute	30 seconds			
Utkattasana			1 minute	30 seconds			
Gomukasana			1 minute	30 seconds			
Meditation – Omkar.			1 minute	30 seconds			
Pranayama – Sitali.			1 minute	30 seconds			
Shavasana			2 minutes		1		
As in previous week	7-9 Weeks	66 min	40 minutes		3	Monday Tuesday Wednesday Thursday Friday & Saturday	5 – 10 Minutes
Sedhupandhasan			1 minutes	30 seconds			
Matsyasana			1 minutes	30 seconds			
Uttanasana			1 minute	30 seconds			
Meditation - Omkar			1 minutes	30 seconds			
Pranayama – Bhastrika.			1 minute	1 minute			
Shavasana			2 minutes		1		
As in previous week	10-12 Weeks	86 min	66 minutes		3	Monday Tuesday Wednesday Thursday Friday & Saturday	5 – 10 Minutes
Paschimottasana			1 minute	30 seconds			
Ushatrasana			1 minute	30 seconds			
Meditation – Omkar.			1 minutes	30 seconds			
Pranayama – ujjayi.			1 minute	30 seconds			
Shavasana							



S. Chidambara Raja

Associate Professor, Department of Physical Education and Sports Sciences, Annamalai University.